

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	X

Pomalyst (pomalidomide)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	1 year

Medications	Quantity Limits
Pomalyst (pomalidomide)	May be subject to quantity limit

APPROVAL CRITERIA

Requests for Pomalyst (pomalidomide) may be approved if the following criteria are met:

- I. Individual has a diagnosis of relapsed/refractory Multiple Myeloma; **AND**
- II. Individual is using in combination with dexamethasone (with or without a proteasome inhibitor or cyclophosphamide); **AND**
- III. Individual has demonstrated disease progression on or within 60 days of completion of the last therapy; **AND**
- IV. Individual has had a trial of at least two prior therapies, including an immunomodulatory agent (such as lenalidomide and proteasome inhibitor) (Label, NCCN 2A);

OR

- V. Individual has a diagnosis of relapsed/refractory Multiple Myeloma; **AND**
- VI. Individual has had a trial of at least two prior therapies, including an immunomodulatory agent (such as lenalidomide) and a proteasome inhibitor; **AND**
- VII. Individual is using in combination with dexamethasone and one of the following (NCCN 2A):
 - A. Daratumumab (Darzalex or Darzalex Faspro); **OR**
 - B. Elotuzumab; **OR**
 - C. Isatuximab-irfc;

OR

- VIII. Individual has a diagnosis of systemic light chain amyloidosis and is using in combination with dexamethasone (NCCN 2A);

OR

- IX. Individual has a diagnosis of AIDS-Related Kaposi Sarcoma (KS); **AND**
- X. Highly active antiretroviral therapy (HAART) has failed to control disease (i.e. stable or increasing KS lesions despite HAART);

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	X

OR

XI. Individual has a diagnosis of Kaposi Sarcoma; **AND**

XII. Individual is HIV-negative.

Note:

Pomalyst (pomalidomide) has black box warnings for embryo-fetal toxicity and venous and arterial thromboembolism. Pomalyst, as a thalidomide analogue, is contraindicated in pregnancy. Thalidomide is a known human teratogen causing severe life-threatening birth defects. Exclude pregnancy with 2 negative pregnancy tests before starting treatment and prevent pregnancy with 2 reliable contraception methods, during and for 4 weeks after treatment. To avoid embryo-fetal exposure, Pomalyst is only available through a restricted distribution program, the POMALYST REMS program. In addition, deep venous thrombosis (DVT), pulmonary embolism (PE), myocardial infarction, and stroke have occurred in patients with MM treated with Pomalyst. Thromboprophylaxis is recommended and should be based on underlying risks.

Key References:

1. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2020. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
2. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: May 28, 2020.
3. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
4. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2020; Updated periodically.
5. Tun HW, Johnston PB, DeAngelis LM, et al. Phase 1 study of pomalidomide and dexamethasone for relapsed/refractory primary CNS or vitreoretinal lymphoma. *Blood* 2018;132:2240-2248.
6. NCCN Clinical Practice Guidelines in Oncology™. © 2019 National Comprehensive Cancer Network, Inc. For additional information visit the NCCN website: <http://www.nccn.org/index.asp>. Accessed on October 10, 2019.
 - a. AIDS-Related Kaposi Sarcoma V2.2020. Revised June 1, 2020.
 - b. Central Nervous System Cancers V2.2019. Revised September 16, 2019.
 - c. Multiple Myeloma V2.2020. Revised October 9, 2019.
 - d. Systemic Light Chain Amyloidosis V1.2019. Revised October 26, 2018.

Federal and state laws or requirements, contract language, and Plan utilization management programs or policies may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.